

02 Dec 2025
Vol. 36, No. 8

For Additional Information:

Dr. Robert Junod, (256) 961-7743

rob.junod@uah.edu

Dr. John Christy, (256) 961-7763

christy@nsstc.uah.edu

Dr. Roy Spencer, (256) 961-7960

spencer@nsstc.uah.edu

Global Temperature Report: November 2025 with Version 6.1

Global climate trend since Dec. 1 1978: +0.16 C per decade

November Temperatures v6.1 (preliminary)

Global composite temp: +0.43°C (+0.77°F) above the seasonal average

Northern Hemisphere: +0.59°C (+1.06 °F) above seasonal average

Southern Hemisphere: +0.27°C (+0.48°F) above seasonal average

Tropics: +0.24°C (+0.44°F) above seasonal average

October Temperatures v6.1 (final)

Global composite temp: +0.53°C (+0.96°F) above the seasonal average

Northern Hemisphere: +0.52°C (+0.93 °F) above seasonal average

Southern Hemisphere: +0.55°C (+0.99°F) above seasonal average

Tropics: +0.24°C (+0.44°F) above seasonal average

Notes on data released Dec 02, 2025

[Please note that we provide these data out of our own initiative, and are only able to produce these updates at times convenient to our working schedules.]

The global mean November temperature departure from the seasonal average dropped slightly to +0.43°C (+0.77°F) from +0.53°C (+0.96°F) in October. The cooling atmosphere across the Southern Hemisphere dominated the decline. The 46+ year trend is currently at +0.156°C/decade but is rounded up to +0.16°C/decade. We estimate the error range of this trend over 46+ years at $\pm 0.03^\circ\text{C/decade}$, which renders the third decimal inconsequential.

As mentioned in last month's report, cool La Niña conditions are officially present in the tropical Pacific. La Niña conditions are expected to persist through December 2025-February 2026, with a transition back to "neutral" conditions likely (61% chance) during January-March 2026. For the latest in the El Niño/La Niña situation, see:

https://www.cpc.ncep.noaa.gov/products/analysis_monitoring/lanina/enso_evolution-status-fcsts-web.pdf

The planet's warmest atmospheric temperature departure in November occurred in central Greenland near Summit Station at +6.01°C (+10.82°F). Other warm areas appeared over most of Canada, the United States, Eastern Europe, Western Russia, Western China, and East Antarctica.

With a reading of -2.76°C (-4.97°F), the coolest departure from average was found in the Southern Ocean south of Africa, near Port Alfred on Possession Island, part of the Subantarctic Crozet Archipelago. Colder than average temperatures were also found in other parts of the Southern Ocean, northern Russia, the Northeast United States, and near Svalbard.

The conterminous US in November was much warmer than average, with temperatures +1.32°C (+2.38°F) above the seasonal mean. The western half of the U.S. experienced the warmest departures from normal, while the Northeast recorded below-normal temperatures. Alaska was slightly cooler, so the 49-state average came in at +1.23°C (+2.21°F) above normal. [We don't include Hawaii in the US results because its land area is less than that of one satellite grid square, so it would have virtually no impact on the overall national results.]

Background notes.

New v6.1 due to termination of NOAA-19 in 2021 and adding METOP-C.

<https://www.drroyspencer.com/2024/11/uah-global-temperature-update-for-october-2024-truncation-of-the-noaa-19-satellite-record/>

New Reference Base Jan 2021 and forward. As noted in the Jan 2021 GTR, the situation comes around every 10 years when the reference period or "30-year normal" that we use to calculate the departures is redefined. With that, we have averaged the absolute temperatures over the period 1991-2020, in accordance with the World Meteorological Organization's guidelines, and use this as the new base period. This allows the anomalies to relate more closely to the experience of the average person, i.e. the climate of the last 30 years. Due to the rising trend of global and regional temperatures, the new normals are a little warmer than before, i.e. the global average temperature for Januaries for 1991-2020 is 0.14 °C warmer than the average for Januaries during 1981-2010. So, the new departures from this now warmer average will appear to be cooler, but this is an artifact of simply applying a new base period. It is important to remember that changes over time periods, such as a trend value or the relative difference of one year to the next, will not change. Think about it this way, all we've done is to take the *entire* time series and shifted it down a little.

To-Do List: There has been a delay in our ability to utilize and merge the new generation of microwave sensors (ATMS) on the NPP and JPSS satellites, but we are renewing our efforts as Dr. Braswell is now focused on this task. The delay is due to the incredibly slow rate at which the data may be accessed. In addition, the current non-drifting satellite operated by the Europeans, MetOP-B, has not yet been adjusted or “neutralized” for its seasonal peculiarities related to its unique equatorial crossing time (0930). While these MetOP-B peculiarities do not affect the long-term global trend, they do introduce error within a particular year in specific locations over land. We have now added MetOP-C to replace the truncated data from NOAA-19.

Dr. Christy and Dr. Roy Spencer, an ESSC principal scientist, use data gathered by advanced microwave sounding units on NOAA, NASA and European satellites to produce temperature readings for almost all regions of the Earth. This includes remote desert, ocean and rain forest areas where reliable climate data are not otherwise available. Dr. Danny Braswell has reconstituted the code which converts the satellite radiances to temperature values and Dr. Rob Junod prepares the monthly reports as of October 2025.

The satellite-based instruments measure the temperature of the atmosphere from the surface up to an altitude of about nine kilometers above sea level. Once the monthly temperature data are collected and processed, they are placed in a "public" computer file for immediate access by atmospheric scientists in the U.S. and abroad.

The complete version 6.1 lower troposphere dataset is available here:

http://www.nsstc.uah.edu/data/msu/v6.1/tlt/uahncdc_lt_6.1.txt

Archived color maps of local temperature anomalies are available on-line at:

<http://nsstc.uah.edu/climate/>

Neither Christy nor Spencer receives any research support or funding from oil, coal or industrial companies or organizations, or from any private or special interest groups. All of their climate research funding comes from federal and state grants or contracts.

NOVEMBER 2025
LAYER = LT LOWER TROPOSPHERE




Figure 1. Lower tropospheric temperature anomalies for September 2025. Contour interval 1.0°C




Figure 2. Bar chart of global monthly lower tropospheric temperature anomalies